

Cough, Cold & Allergy Dosage Chart

We do NOT recommend that children receive cough and cold medicine if they are <u>less than 4 years of age</u>, unless otherwise directed by your child's provider. Please contact our office if you believe cough and cold medicine will benefit your child so we can assist you in administering the proper dosage.

Allegra (Fexofenadine)							
Age	Dosage	Children's Liquid	Children's Tablets	60mg/12hr Tablets	180mg/24hr Tablets		
Under 2 Years		Not Recommended	Not Recommended				
2-6 Years	30mg	1 tsp every 12 hours	1 tablet every 12 hours				
6-12 Years	30mg	1 tsp every 12 hours	1 tablet every 12 hours				
12+ Years	60mg	2 tsp every 12 hours	2 tablets every 12 hours	1 tablet every 12 hours	1 tablet every 24 hours		

Benadryl (Diphenhydramine) May be taken every 4-6 hours					
Weight	Liquid Benadryl 12.5mg/5ml	Chewable 12.5mg	Tablets 25mg		
18-28 lbs	3/4 tsp (3.75ml)				
29-41 lbs	1 tsp (5ml)	1 Chewable			
42-54 lbs	1.5 tsp (7.5ml)				
55-83 lbs	2 tsp (10ml)	2 Chewable	1 Tablet		
84-110 lbs	3 tsp (15ml)	3 Chewable			
110+ lbs	4 tsp (20ml)	4 Chewable	2 Tablets		

Claritin (Loratadine)						
Age	Syrup 1mg/1ml	Chewable Tablets 5mg	Tablets 10mg			
Under 2 Years	Not Recommended					
2-5 Years	1 tsp Once Daily (5ml)	1 Tablet Once Daily				
6+ Years	2 tsp Once Daily (10ml)	2 Tablets Once Daily	1 Tablet Once Daily			

Zyrtec (Cetirizine)						
Age	Syrup 1mg/1ml	Chewable Tablets 5mg	Tablets 10mg			
Under 2 Years	½ tsp Once Daily (2.5ml)					
2-5 Years	½ to 1 tsp (5ml) Once Daily	½ to 1 Tablet Once Daily				
6+ Years	2 tsp Once Daily (10ml)	2 Tablets Once Daily	1 Tablet Once Daily			