



## Acetaminophen/Ibuprofen Dosage Chart

### Acetaminophen (Tylenol<sup>®</sup>)

Give every 4-6 hours as needed. Do not give more than 5 doses in 24 hours.

Weight in pounds (lbs.)	Elixir 1 teaspoon 160 mg/5 ml	Chewable 1 tablet 80 mg	Jr. Strength 1 caplet 160 mg	Reg. Strength 1 tablet 325 mg
6-11 lbs.	1.25 ml	-----	-----	-----
12-17 lbs.	2.5 ml	-----	-----	-----
18-23 lbs.	3.75 ml	1 ½ tablets	-----	-----
24-35 lbs.	5 ml	2 tablets	1 tablet	-----
36-47 lbs.	7.5 ml	3 tablets	1 ½ tablets	-----
48-59 lbs.	10 ml	4 tablets	2 caplets	1 tablet
60-71 lbs.	12.5 ml	5 tablets	2 ½ caplets	1 tablet
72-95 lbs.	15 ml	6 tablets	3 caplets	1 ½ tablets
96+ lbs.	-----	-----	4 caplets	2 tablets

### Ibuprofen (Advil<sup>®</sup>, Motrin<sup>®</sup>)

Give every 6 to 8 hours as needed. Not intended for use by children under 6 months of age. Do not give more than 4 doses in 24 hours.

Weight in pounds (lbs)	Infant Liquid 50 mg/1.25 mL	Children's Liquid 100 mg/5 mL	Children's Chewables 100 mg tablet	Children's 100 mg tablet	Adult 200 mg tablet
12-17 lbs.	1.25 mL	2.5 mL	-----	-----	-----
18-23 lbs.	1.875 mL	4 mL	-----	-----	-----
24-35 lbs.	2.5 mL	5 mL	1 tablet	-----	-----
36-47 lbs.	3.75 mL	7.5 mL	1 ½ tablet	-----	-----
48-59 lbs.	-----	10 mL	2 tablets	2 tablets	1 tablet
60-71 lbs.	-----	12.5 mL	2 ½ tablets	2 tablets	1 tablet
72-95 lbs.	-----	15 mL	3 tablets	3 tablets	1 tablet
96+ lbs.	-----	20 mL	4 tablets	4 tablets	2 tablets